

PENGUIN AMPLITUDES IN HADRONIC B DECAYS: NLO SPECTATOR SCATTERING

SEBASTIAN JÄGER*

Institut für Theoretische Physik E, RWTH Aachen, D-52056 Aachen, Germany

Address after 01 October 2006: *Arnold Sommerfeld Center, Department für Physik,
Ludwig-Maximilians-Universität München, Theresienstraße 37, D-80333 München, Germany*

*E-mail: *jaeger@theorie.physik.uni-muenchen.de*

We present results on the NLO (α_s^2) spectator-scattering corrections to the topological penguin amplitudes for charmless hadronic two-body B -decays in QCD factorization. The corrections can be sizable for the colour-suppressed electroweak penguin amplitudes $\alpha_{4,\text{EW}}^p$ but otherwise are numerically small. Our results explicitly demonstrate factorization at this order. To assess the phenomenological viability of the framework, we consider penguin-to-tree ratios in the penguin-dominated πK system and find agreement to the expected precision (i.e., a power correction).

Keywords: QCD; Factorization; B -meson decays.

1. Introduction

Branching ratios and CP asymmetries in B -decays into two light mesons^a provide access to the flavour structure of the Standard Model (CKM matrix elements) and its possible extensions. Within the Standard Model, the theoretical expressions always involve two terms with a relative weak phase,

$$\mathcal{A}(\bar{B} \rightarrow M_1 M_2) = T_{M_1 M_2} e^{-i\gamma} + P_{M_1 M_2}. \quad (1)$$

Direct CP asymmetries are then governed, besides γ , by the imaginary part $\text{Im } P/T$. A nontrivial theoretical task is to evaluate the strong amplitudes P and T . Integrating out the weak scale by means of the weak effective Hamiltonian, this reduces to the computation of hadronic matrix elements of local operators Q_i , a task currently not feasible on the lattice. Fortunately, at leading power in an expansion in Λ_{QCD}/m_b they obey¹

$$\begin{aligned} &\langle M_1 M_2 | Q_i | \bar{B} \rangle \\ &= F^{B M_1}(0) \int du T_i^{\text{I}}(u) \phi_{M_2}(u) + (M_1 \leftrightarrow M_2) \\ &+ \int d\omega du dv T_i^{\text{II}}(\omega, v, u) \phi_B(\omega) \phi_{M_1}(v) \phi_{M_2}(u). \end{aligned} \quad (2)$$

The hard-scattering kernels $T_i^{\text{I}}, T_i^{\text{II}}$ are perturbatively calculable as series in α_s , while the form factors $F^{B M_1}$ and the light-cone distribution amplitudes (LCDAs) ϕ encapsulate universal nonperturbative properties of the initial- and final-state particles. An important outcome is that all strong phases are contained in the hard kernels. T_i^{I} is currently known to $\mathcal{O}(\alpha_s)$,¹ while the computation of $T_i^{\text{II},2,3,4}$ has recently been completed at $\mathcal{O}(\alpha_s^2)$ by evaluating the one-loop spectator-scattering corrections to the (topological) penguin amplitudes.⁴ As stated above, the latter are crucial for any direct CP asymmetry in the Standard Model. Moreover they are important for the branching fractions particularly of the penguin-dominated $\Delta S = 1$ modes such as $\bar{B} \rightarrow \pi \bar{K}$. Spectator-scattering contributions to their strong phases and effects proportional to the large Wilson coefficient C_1 appear first at this order, similarly to the colour-suppressed tree amplitude considered in Ref. 2.

2. Effective theory and matching

The kernels T_i^{II} receive contributions from two hard scales $\mu_b \sim m_b, \mu_{\text{hc}} \sim \sqrt{m_b \Lambda}$. They

^aWe restrict ourselves to flavour- $SU(3)$ -nonsinglet mesons in this note.

correspond to hard and hard-collinear terms in an expansion by momentum regions.⁵ The result is further factorization $T_i^{\text{II}}(\omega, v, u) = \int dz H_i^{\text{II}}(\mu_b; u, z) J(\mu_{\text{hc}}; \omega, v, z)$.^{6,7} The coefficient functions are conveniently obtained in a two-step matching onto soft-collinear effective theory,^{8,9} $\text{QCD} \rightarrow \text{SCET}_\text{I} \rightarrow \text{SCET}_\text{II}$, integrating out subsequently the hard and hard-collinear scales. The hard coefficients T_i^{I} and H_i^{II} are interpreted as Wilson coefficients in SCET_I and are found by solving the matching equation

$$Q_i = \int dt T_i^{\text{I}}(t) [\bar{\chi}(tn_-) \chi(0)] \left[C_{A0} [\bar{\xi}(0) h_v(0)] - \frac{1}{m_b} \int ds C_{B1}(s) [\bar{\xi}(0) D_{\perp\text{hc}1}(sn_+) h_v(0)] \right] + \frac{1}{m_b} \int dt ds H_i^{\text{II}}(t, s) [\bar{\chi}(tn_-) \chi(0)] \times [\bar{\xi}(0) D_{\perp\text{hc}1}(sn_+) h_v(0)], \quad (3)$$

where the (schematic) rhs involves SCET_I collinear fields for the directions of motion of M_2 ($\chi, \bar{\chi}$) and M_1 ($\xi, A_{\perp\text{hc}1}$) as well as soft fields h_v and $\bar{q}_s$ suitable for interpolating the B -meson, sandwiched between suitable partonic states. The peculiar form of the second bracket in the first convolution is designed to reproduce full-QCD heavy-to-light form factors as in (2). Of interest to us is the second term in (3), which includes all hard interactions of M_2 with the spectator quark. Decoupling properties of SCET_I suggest that its hadronic matrix element factorizes into a light-cone distribution amplitude $\langle M_2 | [\bar{\chi}\chi] | 0 \rangle \propto \phi_{M_2}$ and a nonlocal object $\Xi^{BM_1}(s) = \langle M_1 | [\bar{\xi}(0) D_{\perp\text{hc}1}(sn_+) h_v(0)] | \bar{B} \rangle$. This expectation is indeed confirmed by the finiteness of the convolutions, found in all currently available computations.

Several fermion-line topologies occur in the full-QCD amplitude differing in how the quark fields in Q_i are contracted with the fields interpolating the external states and/or with each other. Each pair (Q_i , topology) contributes to one of a few operators in the effective theory that are dis-

tinguished, besides the chirality of the light fields, only by their flavour content. Their matrix elements define scale- and scheme-independent amplitude coefficients α_i . One-loop spectator-scattering corrections to the colour-allowed and colour-suppressed topological “trees” α_1, α_2 have been computed in Refs. 2, 3, and the corresponding QCD penguin amplitudes α_3^p, α_4^p ($p = u, c$) as well as the electroweak penguin amplitudes $\alpha_{3,\text{EW}}^p, \alpha_{4,\text{EW}}^p$ are given in Ref. 4.

To arrive at the final form (2) one has to perform a second matching step^b corresponding to the matching equation

$$\int d^4x T \left(\mathcal{L}_{\text{SCET}_\text{I}}^{(1)}(x) [\bar{\xi}(0) D_{\perp\text{hc}1}(sn_+) h_v(0)] \right) = \int dw dr J(w, r) [\bar{\xi}(rn_+) \xi(0)] [\bar{q}_s(w n_-) h_v(0)].$$

The jet function J containing the hard-collinear physics appears identically in the factorization formula for form factors¹⁰ and is known to NLO¹¹, while the brackets $[\bar{\xi}\xi]$ and $[\bar{q}_s h_v]$ result in LCDAs for M_1 and the B -meson once hadronic matrix elements are taken.

3. Penguin amplitudes

Some diagrams related to computing the coefficients H_i^{II} relevant to the penguin amplitudes are shown in Fig. 1. In each diagram, the line to the left is the b -quark, the up-going lines are collinear with M_2 , the right-going line (hard-)collinear with M_1 , and the external vector boson is a hard-collinear-1 gluon or photon. Of the first row in the Figure, the left diagram contributes to the (colour-suppressed) electroweak penguin amplitude $\alpha_{4,\text{EW}}^p$ ($p = u$ or c), due to the photon exchanged between quark lines. Tadpole diagrams like the one on the right vanish when summed. On the other hand,

^bAlternatively, one could try to extract $\Xi^{BM_1}(s)$ from experiment.⁷ This is not feasible beyond LO because the full s -dependence is needed.




Fig. 1. Sample diagrams related to the penguin amplitudes. The upper row contributes to $\alpha_{4,EW}^P$ while the lower row introduces isospin breaking in α_4^P .

the diagrams on the second row induce the flavour structure of the QCD penguin amplitude α_4^P due to the exchanged gluon, but the hard-collinear photon implies isospin breaking. Such small electromagnetic corrections to the QCD penguin amplitudes are omitted here.^c When the photons in the upper row are replaced by gluons, contributions to α_4^P arise. There are more diagrams, including ones without quark loops but with insertions of penguin operators from the weak Hamiltonian. With the input parameter ranges given in Ref. 4 we obtain for the leading-power contribution a_4^c to the amplitude α_4^c :

$$\begin{aligned}
 a_4^c(\pi\pi) &= -0.029 \\
 &\quad -[0.002 + 0.001i]_V - [0.001 + 0.007i]_P \\
 &\quad + \left[\frac{r_{sp}}{0.485} \right] \left\{ [0.001]_{LO} + [0.001 + 0.001i]_{HV} \right. \\
 &\quad \left. + [0.000 - 0.000i]_{HP} + [0.001]_{tw3} \right\} \\
 &= -0.028_{-0.003}^{+0.005} + (-0.006_{-0.002}^{+0.003})i. \quad (4)
 \end{aligned}$$

The contributions labeled “V” and “P” originate from one-loop vertex and penguin corrections to T_i^I in the first (form-factor) term in (2), while the terms “HV” and

“HP” denote the newly computed one-loop spectator-scattering corrections. The term “tw3” denotes an estimate of the twist-3 tree-level spectator scattering contribution, which while being a power correction is by convention included a_4^P . The numbers show that the impact of the new corrections is very small. This is somewhat surprising as the large Wilson coefficient C_1 is involved in the “HP” terms. Closer inspection shows a numerical cancellation between diagrams carrying different colour factors, the origin of which is unclear. The corrections to the colour-allowed electroweak penguin amplitude $\alpha_{3,EW}$ are also very small. The colour-suppressed electroweak penguin amplitude $\alpha_{4,EW}$ receives a larger correction. The correction to its leading-power part a_{10} can be $\mathcal{O}(100\%)$ with respect to the $\mathcal{O}(\alpha_s)$ result.⁴ This is because, like the colour-suppressed tree amplitude α_2 , $\alpha_{4,EW}$ is especially sensitive to spectator scattering due to a numerical cancellation between naive factorization and the 1-loop correction to the first term in (2). For the same reason, it is also more sensitive to uncertainties in hadronic input parameters, most importantly the inverse moment λ_B^{-1} of the B -meson light-cone distribution amplitude. Altogether, perturbation theory appears to be well behaved and significant changes in the predictions for branching fractions and CP asymmetries at this time will be due mainly to changes of hadronic input parameters such as form factors and light-cone distribution amplitudes.

4. Phenomenological implications

Because of the smallness of the corrections compared to uncertainties due to hadronic input parameters, we do not give updated numbers for the branching fractions here. Instead we consider a penguin-to-tree ratio, for which part of the nonperturbative uncertainties cancel out, but which nevertheless can be related to experimental data.

^cIncluding them consistently would necessitate taking into account isospin-breaking effects in the form factors, LCDAs, and decay constants, among other complications.




Fig. 2. Comparing the PP penguin amplitude to data. For an explanation, see the text.

Following Ref. 12, Fig. 2 shows the ratio $\hat{\alpha}_4^c(M_1 M_2)/(\alpha_1(\pi\pi) + \alpha_2(\pi\pi))$ for the pseudoscalar-pseudoscalar (PP) final state $M_1 M_2 = \pi\bar{K}$. Here $\hat{\alpha}_4^c(\pi\bar{K}) = a_4^c(\pi\bar{K}) + r_\chi^{\bar{K}} a_6^c(\pi\bar{K}) + \beta_3^c(\pi\bar{K})$, where $r_\chi^{\bar{K}} a_6^c(\pi\bar{K})$ is a numerically large “charming penguin” power correction that factorizes at $\mathcal{O}(\alpha_s)$ and $\beta_3^c(\pi\bar{K})$ models the (within QCD factorization) incomputable penguin annihilation amplitude. The cross shows the theoretical prediction with errors combined in quadrature (the onion-shaped regions are various estimates of the annihilation contribution, the blue one corresponding to the expected magnitude for this power correction). The grey ring and yellow wedge can be inferred from data on $\bar{B} \rightarrow \pi\pi$ and $\bar{B} \rightarrow \pi\bar{K}$ with very little theory input, where the lighter-coloured areas also including a generous uncertainty on modulus and phase of V_{ub} . The wedge opening to the right is disfavoured by data. We observe that theory and experiment, which includes $A_{CP}(\bar{B}^0 \rightarrow \pi^+ K^-)$, agree within errors, which is nontrivial. Some annihilation contribution is needed, but at the level expected for a power correction. It is conceivable that a large one-loop spectator-scattering correction to a_6^c might have a sim-

ilar impact.

In conclusion, factorization works after inclusion of NLO spectator-scattering effects. The corrections are small except for the colour-suppressed electroweak penguin amplitude. The penguin-to-tree ratios relevant to $\Delta S = 1$ decays are consistent with data at the level of a power correction.

Acknowledgements

The speaker thanks the organizers and his collaborator Martin Beneke. This work was supported in part by the DFG Sonderforschungsbereich/Transregio 9 “Computergestützte Theoretische Teilchenphysik”.

References

1. M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, *Phys. Rev. Lett.* **83**, 1914 (1999); *Nucl. Phys.* **B591**, 313 (2000); *Nucl. Phys.* **B606**, 245 (2001).
2. M. Beneke and S. Jäger, *Nucl. Phys.* **B751**, 160 (2006).
3. N. Kivel, hep-ph/0608291.
4. M. Beneke and S. Jäger, hep-ph/0610322.
5. M. Beneke and V. A. Smirnov, *Nucl. Phys.* **B522**, 321 (1998).
6. J. Chay and C. Kim, *Nucl. Phys.* **B680**, 302 (2004).
7. C. W. Bauer, D. Pirjol, I. Z. Rothstein and I. W. Stewart, *Phys. Rev.* **D70**, 054015 (2004).
8. C. W. Bauer, S. Fleming, D. Pirjol and I. W. Stewart, *Phys. Rev.* **D63**, 114020 (2001); C. W. Bauer, D. Pirjol and I. W. Stewart, *Phys. Rev.* **D65**, 054022 (2002).
9. M. Beneke, A. P. Chapovsky, M. Diehl and T. Feldmann, *Nucl. Phys.* **B643**, 431 (2002); M. Beneke and T. Feldmann, *Phys. Lett.* **B553**, 267 (2003).
10. M. Beneke and Th. Feldmann, *Nucl. Phys.* **B685**, 249 (2004).
11. R. J. Hill, T. Becher, S. J. Lee and M. Neubert, *JHEP* **0407**, 081 (2004); T. Becher and R. J. Hill, *JHEP* **0410**, 055 (2004); M. Beneke and D. Yang, *Nucl. Phys.* **B736**, 34 (2006); G. G. Kirilin, hep-ph/0508235.
12. M. Beneke and M. Neubert, *Nucl. Phys.* **B675**, 333 (2003).